

PubMed

**Format:** Abstract

Full text links



*J Sex Med.* 2017 Jul;14(7):891-897. doi: 10.1016/j.jsxm.2017.05.012.

## Low-Intensity Shockwave Therapy Improves Hemodynamic Parameters in Patients With Vasculogenic Erectile Dysfunction: A Triplex Ultrasonography-Based Sham-Controlled Trial.

[Kalyvianakis D<sup>1</sup>](#), [Hatzichristou D<sup>2</sup>](#).

### Author information

### Erratum in

[Erratum.](#) [*J Sex Med.* 2018]

### Abstract

**BACKGROUND:** Although several reports have documented the subjective improvement of erectile function after low-intensity extracorporeal shockwave therapy (LI-ESWT) in patients with vasculogenic erectile dysfunction (ED), objective assessment data of penile hemodynamics are lacking.

**AIM:** To assess penile hemodynamics before and 3 months after LI-ESWT in a group of patients with documented vasculogenic ED.

**METHODS:** This was a double-blinded, randomized, sham-controlled trial. Forty-six patients with ED were randomized; 30 underwent LI-ESWT and 16 had a sham procedure in double-blinded fashion. All patients underwent penile triplex ultrasonography by the same investigator immediately before and 3 months after treatment. Patient demographics, International Index of Erectile Function erectile function domain (IIEF-ED) score, and minimal clinically important difference were assessed at baseline and 1, 3, 6, 9, and 12 months after treatment.

**OUTCOMES:** Changes in peak systolic velocity and resistance index as measured by triplex ultrasonography at baseline and 3 months after treatment were the main outcomes of the study. Secondary outcomes were changes in the IIEF-ED score from baseline to 1, 3, 6, 9, and 12 months after treatment and the percentage of patients reaching a minimal clinically important difference during the same period for the two groups.

**RESULTS:** IIEF-ED minimal clinically important differences for the active vs sham group were observed for 56.7% vs 12.5% ( $P = .005$ ) at 1 month, 56.7% vs 12.5% ( $P = .003$ ) at 3 months, 63.3% vs 18.8% ( $P = .006$ ) at 6 months, 66.7% vs 31.3% ( $P = .022$ ) at 9 months, and 75% vs 25% ( $P = .008$ ) at 12 months. Mean peak systolic velocity increased by 4.5 and 0.6 cm/s in the LI-ESWT and sham groups, respectively ( $P < .001$ ).

**CLINICAL IMPLICATIONS:** Such results offer objective and subjective documentation of the value of this novel treatment modality for men with vasculogenic ED.

**STRENGTHS AND LIMITATIONS:** Strengths include the prospective, randomized, sham-controlled type of study and the assessment of penile hemodynamics.

Limitations include the small sample and strict inclusion criteria that do not reflect everyday clinical practice.

**CONCLUSION:** The present study confirms the beneficial effect of LI-ESWT on penile hemodynamics and the beneficial effect of this treatment up to 12 months. Kalyvianakis D, Hatzichristou D. Low-Intensity Shockwave Therapy Improves Hemodynamic Parameters in Patients With Vasculogenic Erectile Dysfunction: A Triplex Ultrasonography-Based Sham-Controlled Trial. *J Sex Med* 2017;14:891-897.

Copyright © 2017 International Society for Sexual Medicine. Published by Elsevier Inc. All rights reserved.

**KEYWORDS:** Erectile Dysfunction; Low-Intensity Shockwave Therapy; Peak Systolic Velocity; Penile Doppler

PMID: 28673433 DOI: [10.1016/j.jsxm.2017.05.012](https://doi.org/10.1016/j.jsxm.2017.05.012)

[Indexed for MEDLINE]

---

**Publication type, MeSH terms**

---

**LinkOut - more resources**